

Perturbative continued-fraction method for weakly interacting Brownian spins and dipoles

J. L. García-Palacios

Dept. de Teoría y Simulación de Sistemas Complejos,*

Instituto de Ciencia de Materiales de Aragón, C.S.I.C. – Universidad de Zaragoza, E-50009 Zaragoza, Spain

The continued-fraction (CF) method was developed systematically by Risken and co-workers to solve problems of arbitrary fluctuations in nonlinear systems¹. However, this efficient technique is limited to problems with a few variables,² which in practice means systems of noninteracting entities (particles, spins, etc.) In this communication, we illustrate how to extend the CF method to weakly *coupled* systems with the problem of relaxation in classical spins.

The Fokker–Planck equation governing the $T \neq 0$ dynamics of a classical spin $\vec{s}$ is given by³

$$2\tau_D \frac{\partial W}{\partial t} = -\frac{\partial}{\partial \vec{s}} \cdot \left\{ \frac{1}{\lambda} \vec{s} \wedge \vec{B} - \vec{s} \wedge \left[\vec{s} \wedge \left(\vec{B} - \frac{\partial}{\partial \vec{s}} \right) \right] \right\} W,$$

where λ is the dissipation parameter, $\vec{B} = -(\partial \mathcal{H} / \partial \vec{s})$ the effective field, and $\tau_D \propto T^{-1}$ the free diffusion time. If we write this equation as $\partial W / \partial t = \mathcal{L}_{\text{FP}} W$, the dynamical equation for any average $\langle f \rangle$ is given in terms of the adjoint Fokker–Planck operator by $d\langle f \rangle / dt = \langle \mathcal{L}_{\text{FP}}^\dagger f \rangle$.

However, the underlying Landau–Lifshitz damping term $-\lambda \vec{s} \wedge (\vec{s} \wedge \vec{B})$ is nonlinear in $\vec{s}$ and couples the equations for the moments. To handle the infinite systems of coupled equations it is convenient to introduce the spherical harmonics $X_l^m = e^{im\varphi} P_l^m(s_z)$, and take advantage of their recurrence relations. Writing $\mathcal{L}_{\text{FP}}^\dagger$ in terms of the rotation operator $\vec{\mathcal{J}} = -i \vec{s} \wedge (\partial / \partial \vec{s})$

$$\mathcal{L}_{\text{FP}}^\dagger = -\frac{i}{\lambda} \vec{B} \cdot [\vec{\mathcal{J}} + \lambda(\vec{s} \wedge \vec{\mathcal{J}})] - \vec{\mathcal{J}}^2, \quad (1)$$

we get the equations for the X_l^m in the form of a recurrence relation that can be solved by CF methods.

When the spins *interact*, the structure of $\mathcal{L}_{\text{FP}}^\dagger$ is the same with the field created by the other spins $\vec{b}_d$ included in $\vec{B}$. This allows to separate the free evolution and interaction terms, $\mathcal{L}_{\text{FP}}^\dagger = \mathcal{L}_{\text{FP},0}^\dagger + \vec{b}_d \cdot \vec{\mathcal{J}}_{\text{LL}}$, where $\vec{\mathcal{J}}_{\text{LL}}$ is the operator multiplying $\vec{B}$ in Eq. (1). For weak coupling, we can expand the averages in the coupling parameter $\langle f \rangle = \langle f \rangle_0 + \langle f \rangle_1 + \dots$, getting a hierarchy of equations

$$\begin{aligned} d\langle f \rangle_0 / dt &= \langle \mathcal{L}_{\text{FP},0}^\dagger f \rangle_0 \\ d\langle f \rangle_1 / dt &= \langle \mathcal{L}_{\text{FP},0}^\dagger f \rangle_1 + \langle \vec{b}_d \cdot \vec{\mathcal{J}}_{\text{LL}} f \rangle_0 \\ d\langle f \rangle_2 / dt &= \langle \mathcal{L}_{\text{FP},0}^\dagger f \rangle_2 + \langle \vec{b}_d \cdot \vec{\mathcal{J}}_{\text{LL}} f \rangle_1 \\ &\vdots \end{aligned}$$

To illustrate our procedure, let us denote by X_ℓ^m and Y_p^q the spherical harmonics of two of the spins, and consider the equation for X_ℓ^m (i.e., $f = X_\ell^m$ above). Since the field created by the second spin depends on Y_p^q , one

needs $\langle X_\ell^m Y_p^q \rangle_i$, $i = 0, 1$, which could be obtained writing the above equations also for $f = X_\ell^m Y_p^q$. This leads to another complication, since $\vec{b}_d \cdot \vec{\mathcal{J}}_{\text{LL}} f$ then involves products of three spherical harmonics. However, $Y_1^\mu Y_p^q$ can be expanded as a linear combination of the Y_p^q 's, restoring a structure solvable by CF methods.

These manipulations result in a complicated algorithm, so it is convenient to check it out against some solvable case. Fortunately, for freely rotating spins the problem reduces to that of electric dipoles, where the perturbative calculation can be done analytically, yielding⁴

$$\frac{\alpha(\omega)}{\alpha_0(\omega)} = 1 - \frac{\alpha^2 \varrho^2}{1 + i\omega\tau_D} \left\{ \mathcal{R} \left[1 + \frac{i\omega\tau_D}{2(4 + i\omega\tau_D)} \right] - \frac{\mathcal{S}}{1 + i\omega\tau_D} \right\}$$

where $\alpha_0(\omega)$ and α ($\propto T^{-1}$) are the dynamic and static polarisabilities of noninteracting dipoles, ϱ is the dipole density, and $\mathcal{R}$ and $\mathcal{S}$ are certain “lattice sums” of the interaction tensor. Figure 1 shows that our perturbative CF approach gives accurately the analytical results.




FIG. 1. Cole-Cole plots of the dielectric polarisability (imaginary vs. real part) for various couplings ϱ . Lines: Zwanzig formula; crosses: perturbative CF results.

Therefore, this approach extends the applicability of the efficient CF techniques to (weakly) interacting systems, and allows one to handle problems involving *driven* systems, evolving in *nonlinear potentials*, *coupled*, and subjected to *dissipation* and *fluctuations*.

The author acknowledges financial support from DGES (PB98-1592) and F. Falo for useful discussions.

* wzar.unizar.es/acad/fac/cie/cond-mat/T/theory.html

¹ H. Risken, *The Fokker–Planck Equation* (Springer, 1989).

² P. Hänggi *et al.*, Rev. Mod. Phys., **62**, 251 (1990)

³ W. F. Brown, Jr., Phys. Rev. **130**, 1677 (1963).

⁴ R. Zwanzig, J. Chem. Phys. **38**, 2766 (1963)